

Analysis of Beta-Casein Gene (CSN2) Polymorphism in Different Breeds of Cattle

Martina Miluchová*, Michal Gábor, Anna Trakovická

Slovak University of Agriculture in Nitra, 949 76 – Nitra, Tr. A Hlinku, 2, Slovakia

Abstract

The goal of work was identification of β - casein gene polymorphism in different breeds of cow. The beta - casein constitutes up to 45 % of the casein of bovine milk. The most common forms of beta-casein in dairy cattle breeds are A1 and A2, while B is less common. The β -casein A1 variant was associated with the incidence of diabetes mellitus 1st type, coronary heart disease and autism. The A2 variant reduces serum cholesterol. The material involved 287 cows (Simmental breed – 111 cows, Pinzgau breed – 89 cows, Holstein breed – 87 cows). Bovine genomic DNA was extracted from whole blood by using commercial kit and used in order to estimate β - casein genotypes by means of PCR-RFLP method. In the populations included in the study were detected all three genotypes – homozygote genotype A1A1, heterozygote genotype A1A2 and homozygote genotype A2A2 with frequencies 0.1261, 0.3333 and 0.5405 in Simmental breed; 0.1379, 0.4598 and 0.4023 in Holstein breed, 0.3034, 0.5168 and 0.1798 in Pinzgau breed. In population of Simmental breed and Holstein breed was higher frequency of allele A2 (0.7072 and 0.6322). In opposite, in population of Pinzgau breed was present higher frequency of the allele A1 (0.5618).

Keywords: beta-casein, cattle, PCR-RFLP

1. Introduction

One of the primary functions of milk is to protect the health of a newborn mammal. Milk contains many peptides and proteins. Milk proteins were classified in two groups: caseins constitute about 80 % of the protein content of milk; the remaining about 20 % of milk protein content consists of whey proteins [1-3].

The beta - casein (CSN2) constitutes up to 45 % of the casein of bovine milk. CSN2 is localised in bovine chromosome 6 [4]. The primary sequence of β -casein has been reported by Ribadeau-Dumas *et al.* (1972) [5]. Single - polypeptide chain of this protein containing 209 residues with molecular weight of 23983.

The most common forms of beta-casein in dairy cattle breeds are A1 and A2, while B is less

common [6-8]. The original beta-casein protein in bovine milk was A2. A1 beta-casein is a consequence of a mutation [9]. The β - casein A1 and B variants differ from the A2 variant at position 67 where a histidine (codon CAT) replaces a proline (codon CCT). In addition, the B variant differs from the A1 variant in a substitution of argine for serine at position 122. Importantly, it is the change to histidine at position 67 that has the potential to result in cleavage occurring upon digestion and a bioactive peptide, beta-casomorphin potentially being liberated [10-12]. Beta-casomorphin (BCM7) have also been suggested to show many immunological activities like chronic inflammatory responses, such as allergy, mucin production, lymphocyte proliferation, skin reactions [13]. Human milk, goat milk, sheep milk and other species are „A2-like“ with proline at the equivalent position [14, 15].

The β -casein A1 variant was associated with the incidence of diabetes mellitus 1st type, coronary

* Corresponding author: Martina Miluchová
Tel: 00421-6414290, martina.miluchova@centrum.sk

heart disease and autism [16]. The A2 variant reduces serum cholesterol [17].

The goal of work was identification of β - casein gene polymorphism in different breeds of cow.

2. Materials and methods

The material involved 287 cows (Slovak spotted breed – 111 cows, Pinzgau breed – 89 cows, Holstein breed – 87 cows). Bovine genomic DNA was extracted from whole blood by using commercial kit and used in order to estimate β - casein genotypes by means of PCR-RFLP method. DNA primers described by McLachlan (2006) [18] were used to PCR amplification: forward primer 5'- CCT TCT TTC CAG GAT GAA CTC CAG G- 3' and reverse primer 5' - GAG TAA GAG GAG GGA TGT TTT GTG GGA GGC TCT- 3'.

The reaction mixture in the total volume 10 μ l containing 50 ng DNA, 1.5 U Taq polymerase (Fermentas), 1X PCR buffer (750 mM Tris-HCl, pH 8.8, 200 mM (NH₄)₂SO₄, 0.1% Tween 20), 6 mM MgCl₂, 200 μ M dNTP, 5 pM of each primer. The following amplification parameters were applied: 95°C for 5 minutes followed by 30 cycles: 95 °C for 10 seconds, 58 °C for 30 seconds, 72 °C for 30 seconds. The reaction was completed by the final extension: 72 °C for 5 minutes.

The PCR products of 121 bp were digested with 5 units of the *Dde*I restriction enzyme (Fermentas). Restriction digestion fragments were loaded on 3 % agarose gel (Invitrogen) containing GelRed™ (Biotium) in 1 \times SB buffer [19] at 180 V for 15 minutes and the gel were analyzed in the UV rays and the documentary system Olympus C-7070 were used to record the results.

3. Results and discussion

*Dde*I digestion of the PCR product was analyzed by 3% agarose-gel electrophoresis. Allele A1 produced 121 bp fragment, and allele A2 produced a 86 bp and 35 bp fragments as the PCR-RFLP.




Figure 1. Representatively results of analysis PCR-RFLP for *CSN2* gene by *Dde*I on 3 % agarose gel.

1,8 – marker 100 bp DNA Ladder (Fermentas);
2,5 – genotype A1A2 (121 bp, 86 bp, 35 bp); 3 –
genotype A1A1(121bp); 4,6,7 – genotype A2A2
(86 bp, 35 bp)

In the populations included in the study were detected all three genotypes. In the population of Slovak Spotted breed we detected homozygote genotype A1A1 (14 cows), heterozygote genotype A1A2 (37 cows) and homozygote genotype A2A2 (60 cows). In the total population of cattle homozygotes A2A2 – 0.5405 were the most frequent, while homozygotes A1A1 – 0.1261 were the least frequent ones. This suggests a superiority of allele A2 (0.7072).

In Holstein breed was observed homozygote genotype A1A1 (12 cows), heterozygote genotype A1A2 (40 cows) and homozygote genotype A2A2 (35 cows) with frequencies 0.1379, 0.4598 and 0.4023, subsequently. In Pinzgau breed was detected homozygote genotype A1A1 (27 cows), heterozygote genotype A1A2 (46 cows) and homozygote genotype A2A2 (16 cows) with frequency 0.3034, 0.5168 and 0.1798, subsequently. . In population of Slovak spotted breed and Holstein breed was higher frequency of allele A2 (0.7072 and 0.6322). In opposite, in population of Pinzgau breed was present higher frequency of the allele A1 (0.5618). Genetic equilibrium of analysed populations was evaluated on the base χ^2 -test. In the populations included in the study non-significant differences in frequencies of genotypes were found.

Detailed genotype and gene frequencies per breed are presented in Table 1.

Table 1. Genotype and allele frequencies of CSN2 gene in different breeds of cow

BREED	COWS	GENOTYPE FREQUENCIES			ALLELE FREQUENCIES		X ² d.f.=2	P
		A1A1	A1A2	A2A2	A1	A2		
Slovak Spotted breed	111	0.1261	0.3333	0.5406	0.2928	0.7072	4.232	0.1205
Pinzgau breed	89	0.3034	0.5168	0.1798	0.5618	0.4382	0.22	0.896
Holstein breed	87	0.1379	0.4598	0.4023	0.3678	0.6322	0.011	0.9945
TOTAL	287							

Table 2. Effectiveness of alleles for CSN2 gene in different breeds of cow

Breed	Locus	Alleles	H _{obs}	H _e	PIC	E	ENA	V%
Slovak Spotted breed	CSN2	A1;A2	0.3333	0.4142	0.3285	0.5858	1.7071	41.80
Pinzgau breed			0.5168	0.4924	0.3712	0.5076	1.9701	49.78
Holstein breed			0.4598	0.4650	0.3568	0.535	1.8692	47.04

The expected homozygosity for gene CSN2 is in all populations stated a slight increase in homozygosity (Slovak Spotted breed – 0.5858, Pinzgau breed – 0.5076, Holstein breed – 0.535). This caused a slight decrease in the level of possible variability realization (Slovak Spotted breed – 41.80%, Pinzgau breed – 49.78%, Holstein breed – 47.04%), which corresponds to the effective number of alleles (Slovak Spotted breed – 1.7071, Pinzgau breed – 1.9701, Holstein breed – 1.8692).

Frequencies of A2 allele in population Slovak Spotted breed and Holstein breed were similar to those of CSN2 gene as reported by Beja-Pereira *et al.* (2003) [20] for Pinzgau cattle. The predominance of CSN2 A2 allele (0.764) detected Caroli *et al.* (2008) [21] in population of Carora cattle. Manga *et al.* (2006) [22] presented lower frequency of the allele A1 in population of Czech Spotted and Czech Holstein breed. In population of Pinzgau breed we detected slight superiority of allele A1. The higher frequency of the allele A1 was reported by Bech *et al.* (1990) [23] for Black-and-White breed and Ehrmann *et al.* (1997) [24] for Red-and-White breed. Ikonen *et al.* (1997) [25] reported slight superiority of allele A1 for Ayrshire breed and Hanusová *et al.* (2010) [26] for Holstein bulls.

4. Conclusions

It may be concluded that Slovak spotted breed and Holstein breed exhibit a superiority of allele A2 (0.7072) which does not produce BCM-7 and thus is safe for human consumption.

Acknowledgements

This work has been supported by the grants: The Slovak Research and Development Agency under the contract No. LPP-0220-09 and No. APVV-0636-11 and Excellence Centre for Agrobiodiversity and Benefit project (ECOVA, ITMS: 26220120015) implemented under the Operational Programme Research and Development financed by European Fund for Regional Development.

References

- Bagnicka, E., Strzałkowska, N., Józwick, A., Krzyżewski, J., Horbańczuk, J.O., and Zwierzchowski, L., Defensins in farm animals, their expression and polymorphism, *Acta Biochim. Pol.*, 2010, 57, 487-497
- Gálik, B., Šimko, M., Juráček, M., Bíro, D., Horniaková, E., Rolinec, M., Pastierik, O., Kolesárová, A., Maiorano, G., Gambacorta, M., Tavaniello, S., and Bednarczyk, M., *Biotechnology and animal food quality*, SPU, Nitra, 2011, 130 p. ISBN 978-80-552-0708-7
- Szwajkowska, M., Wolanciuk, A., Barłowska, J., Król, J., and Litwińczuk, Z., Bovine milk proteins as the source of bioactive peptides influencing the consumers' immune system, *Anim. Sci. Pap. Rep.*, 2011, 29, 269-280
- Ferretti, L., Leone, P., and Sgaramella, U., Long range restriction analysis of the bovine casein gene, *Nuc. Acids. Res.*, 1990, 18, 6829-6833
- Ribadeau-Dumas, B., Brignon, G., Grosclaude, F., and Mercier, J.C., Structure primaire de la caséine β bovine. Séquence complète, *Eur. J. Biochem.*, 1972, 25, 505-514
- Farrell, H.M. Jr., Jimenez-Florez, R., Bleck, G.T., Brown, E.M., Butler, J.E., Creamer, L.K., Hicks, C.L., Hollar, C.M., Ng-Kwai-Hang, K.F., and Swaisgood, G.D., 2003, *Casein: Chemistry and Biochemistry*, CRC Press, Boca Raton, FL, USA, 2003, 1-10

- H.E., Nomenclature of the proteins of cows' milk – sixth revision, *J. Dairy Sci.*, 2004, 87, 1641–1674
7. Kamiński, S., Cieślińska, A., and Kostyra, E., Polymorphism of bovine beta-casein and its potential effect on human health, *J. Appl. Genet.*, 2007, 48, 189–198
8. Keating, A.F., Smith, T.J., Ross, R.P., and Cairns, M.T., A note on the evaluation of a beta-casein variant in bovine breeds by allele-specific PCR and relevance to β -casomorphin, *Irish J. Agr. Food Res.*, 2008, 47, 99–104
9. Ng-Kwai-Hang, K.F., and Grosclaude, F., Genetic polymorphism of milk proteins, In: Fox PFaM, P.L.H editor. *Advanced Dairy Chemistry: Volume 1: Proteins Parts A&B*. New York: Kluwer Academic/Plenum Publishers, 2002, pp. 739-816
10. Stewart, A.F., Bonsing, J., Beattie, C.W., Shah, F., Willis, I.M., and Mackinley, A.G., Complete nucleotide sequences of bovine α 2- and β -casein cDNAs: comparison with related sequences in other species, *Mol. Biol. Evol.*, 1987, 4, 231–241
11. Damiani, G., Pilla, F., Leone, P., and Caccio, S., Direct sequencing and bidirectional allele specific polymerase chain reaction of the bovine β -casein B variant, *Anim. Genet.*, 1992, 23, 561–566
12. Lien, S., Alestrom, P., Klungland, H., and Rogne, S. Detection of multiple β -casein alleles by amplification created restriction sites (ACRS), *Anim. Genet.*, 1992, 23, 333–338
13. Haq, M. R., Kapila, R., Shandilya, U. K., and Kapila, S. Impact of milk derived β -casomorphins on physiological functions and trends in research: A Review, *Internat. J. Food Prop.*, 2014, doi: 10.1080/10942912.2012.712077
14. Lonnerdal, B., Bergstrom, S., Andersson, Y., Hjalmarsson, K., Sundqvist, A.K., Hernell, O., Cloning and sequencing of a cDNA encoding human milk beta-casein, *FEBS Lett.*, 1990, 269, 153-6
15. Provot, C., Persuy, M.A., and Mercier, J.C., Complete nucleotide sequence of ovine beta-casein cDNA: inter-species comparison, *Biochimie*, 1989, 71, 827-32
16. Elliott, R.B., Reddy, S.N., Bibby, N.J., and Kida, K., Dietary prevention of diabetes in the non-obese diabetic mouse, *Diabetologia*, 1988, 31, 62–64.
17. Panicke, L., Freyer, G., Erhardt, G., Effects of milk protein genotypes on milk production traits, 48th Annual Meeting of the European Association for Animal Produktion am 25.-28.08.1997 in Vienna/Austria, 1997
18. McLachlan, C.N., Breeding and milking cows for milk free of β -casein A1, United States Patent 7094949, 2006
19. Brody, R.J., and Kern, S.E., Sodium boric acid: a Tris-free, cooler conductive medium for DNA electrophoresis, *Biotechnique*, 2004, 36, 214-216
20. Beja-Pereira, A., Luikart, G., England, P.R., Bradley, D.G., Jann, O.C., Bertorelle, G., Chamberlain, A.T., Nunes, T.P., Metodiev, S., Ferrand, N., and Erhardt, G., Gene-culture coevolution between cattle milk protein genes and human lactase genes, *Nat. Genet.*, 2003, 35, 311 – 313
21. Caroli, A., Chessa, S., Chiatti, F., Rignanese, D., Meléndez, B., Rizzi, R., and Ceriotti, G., Short communication: Carora cattle show high variability in alpha(s1)-casein, *J. Dairy Sci.*, 2008, 91, 354-9
22. Manga, I., Říha, J., and Dvořák, J., Comparison of influence markers CSN3 and CSN2 on milk performance traits in czech spotted and holstein cattle tested at first, fifth and higher lactation, *Acta fytotech. et zootech.*, 2006, 9, 13-15
23. Bech, A.M., and Kristiansen, K.R., Milk protein polymorphism in Danish dairy cattle and the influence of genetic variants on milk yield, *J. Dairy Res.*, 1990, 57, 53–62.
24. Ehrmann, S., Bartenschlager, H., and Geldermann, H., Quantification of gene effects on single milk proteins in selected groups of dairy cows, *J. Anim. Breed. Genet.*, 1997, 114, 121–132.
25. Ikonen, T., Ojala, M., and Ruottinen, O., Effects of beta and kappa-casein genotypes on first lactation milk production traits in Finnish Ayrshire cows. Milk protein polymorphism, *Int Dairy Fed.*, 1997, 47–53
26. Hanusová, E., Huba, J., Oravcová, M., Polák, P., and Vrtková, I., Genetic variants of beta-casein in Holstein Dairy Cattle in Slovakia, *Slovak. J. Anim. Sci.*, 2010, 43, 63 – 66